

FURTHER READING

Chapter 1

Connor JT. Military medicine, its technology and transition in the mid-twentieth century. *Mil Med.* 2004;169:ix–xi.

Silberman S. The painful truth. *Wired.* 2005;2(13):112–123.

Chapter 3

Buckenmaier CC III, Bleckner LL. Anaesthetic agents for advanced regional anaesthesia: a North American perspective. *Drugs.* 2005;65:745–759.

Tetzlaff JE. The pharmacology of local anesthetics. *Anesthesiol Clin North America.* 2000;18:217–233.

Chapter 4

Dabu A, Chan V. *A Practical Guide to Ultrasound Imaging for Peripheral Nerve Blocks.* Toronto, Canada: University of Toronto; 2004.

De AJ, Sala-Blanch X. Peripheral nerve stimulation in the practice of brachial plexus anesthesia: a review. *Reg Anesth Pain Med.* 2001;26:478–483.

Grau T. Ultrasonography in the current practice of regional anaesthesia. *Best Pract Res Clin Anaesthesiol.* 2005;19:175–200.

Marhofer P, Greher M, Kapral S. Ultrasound guidance in regional anaesthesia. *Br J Anaesth.* 2005;94:7–17.

Chapter 5

Evans H, Steele SM, Nielsen KC, Tucker MS, Klein SM. Peripheral nerve blocks and continuous catheter techniques. *Anesthesiol Clin North America.* 2005;23:141–162.

Lanz E, Theiss D, Jankovic D. The extent of blockade following various techniques of brachial plexus block. *Anesth Analg.* 1983;62:55–58.

Neal JM, Hebl JR, Gerancher JC, Hogan QH. Brachial plexus anesthesia: essentials of our current understanding. *Reg Anesth Pain Med.* 2002;27:402–428.

Chapter 6

Pandit JJ, Bree S, Dillon P, Elcock D, McLaren ID, Crider B. A comparison of superficial versus combined (superficial and deep) cervical plexus block for carotid endarterectomy: a prospective, randomized study. *Anesth Analg.* 2000;91:781–786.

Pandit JJ, Dutta D, Morris JF. Spread of injectate with superficial cervical plexus block in humans: an anatomical study. *Br J Anaesth.* 2003;91:733–735.

Winnie AP, Ramamurthy S, Durrani Z, Radonjic R. Interscalene cervical plexus block: a single-injection technic. *Anesth Analg.* 1975;54:370–375.

Chapter 7

Chan VW: Applying ultrasound imaging to interscalene brachial plexus block. *Reg Anesth Pain Med.* 2003;28:340–343.

Silverstein WB, Saiyed MU, Brown AR. Interscalene block with a nerve stimulator: a deltoid motor response is a satisfactory endpoint for successful block. *Reg Anesth Pain Med.* 2000;25:356–359.

Urmey WF, McDonald M. Hemidiaphragmatic paresis during interscalene brachial plexus block: effects on pulmonary function and chest wall mechanics. *Anesth Analg.* 1992;74:352–357.

Winnie AP. Interscalene brachial plexus block. *Anesth Analg.* 1970;49:455–466.

Chapter 8

Chan VW, Perlas A, Rawson R, Odukoya O. Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg.* 2003;97:1514–1517.

Franco CD, Vieira ZE. 1,001 subclavian perivascular brachial plexus blocks: success with a nerve stimulator. *Reg Anesth Pain Med.* 2000;25:41–46.

Neal JM, Moore JM, Kopacz DJ, Liu SS, Kramer DJ, Plorde JJ. Quantitative analysis of respiratory, motor, and sensory function after supraclavicular block. *Anesth Analg.* 1998;86:1239–1244.

Winnie AP, Collins VJ. The subclavian perivascular technique of brachial plexus anesthesia. *Anesthesiology.* 1964;25:353–363.

Chapter 9

Lecamwasam H, Mayfield J, Rosow L, Chang Y, Carter C, Rosow C. Stimulation of the posterior cord predicts successful infraclavicular block. *Anesth Analg.* 2006;102:1564–1568.

Ootaki C, Hayashi H, Amano M. Ultrasound-guided infraclavicular brachial plexus block: an alternative technique to anatomical landmark-guided approaches. *Reg Anesth Pain Med.* 2000;25:600–604.

Rodriguez J, Barcena M, Rodriguez V, Aneiros F, Alvarez J. Infraclavicular brachial plexus block effects on respiratory function and extent of the block. *Reg Anesth Pain Med.* 1998;23:564–568.

Wilson JL, Brown DL, Wong GY, Ehman RL, Cahill DR: Infraclavicular brachial plexus block: parasagittal anatomy important to the coracoid technique. *Anesth Analg.* 1998;87:870–873.

Chapter 10

Coventry DM, Barker KF, Thomson M. Comparison of two neurostimulation techniques for axillary brachial plexus blockade. *Br J Anaesth.* 2001;86:80–83.

Retzl G, Kapral S, Greher M, Mauritz W. Ultrasonographic findings of the axillary part of the brachial plexus. *Anesth Analg.* 2001;92:1271–1275.

Spence BC, Sites BD, Beach ML. Ultrasound-guided musculocutaneous nerve block: a description of a novel technique. *Reg Anesth Pain Med*. 2005;30:198–201.

Chapter 11

Dilger JA, Wells RE, Jr. The use of peripheral nerve blocks at the elbow for carpal tunnel release. *J Clin Anesth*. 2005;17:621–623.

McCartney CJ, Xu D, Constantinescu C, Abbas S, Chan VW. Ultrasound examination of peripheral nerves in the forearm. *Reg Anesth Pain Med*. 2007;32:434–439.

Chapter 12

Karmakar MK. Thoracic paravertebral block. *Anesthesiology*. 2001;95:771–780.

Klein SM, Bergh A, Steele SM, Georgiade GS, Greengrass RA. Thoracic paravertebral block for breast surgery. *Anesth Analg*. 2000;90:1402–1405.

Naja MZ, Gustafsson AC, Ziade MF, et al. Distance between the skin and the thoracic paravertebral space. *Anaesthesia*. 2005;60:680–684.

Naja MZ, Ziade MF, El Rajab R, El Tayara K, Lonnqvist PA. Varying anatomical injection points within the thoracic paravertebral space: effect on spread of solution and nerve blockade. *Anaesthesia*. 2004;59:459–463.

Chapter 13

Enneking FK, Chan V, Greger J, Hadzic A, Lang SA, Horlocker TT. Lower-extremity peripheral nerve blockade: essentials of our current understanding. *Reg Anesth Pain Med*. 2005;30:4–35.

Evans H, Steele SM, Nielsen KC, Tucker MS, Klein SM. Peripheral nerve blocks and continuous catheter techniques. *Anesthesiol Clin North America*. 2005;23:141–162.

Chapter 14

Awad IT, Duggan EM. Posterior lumbar plexus block: anatomy, approaches, and techniques. *Reg Anesth Pain Med*. 2005;30:143–149.

Capdevila X, Coimbra C, Choquet O. Approaches to the lumbar plexus: success, risks, and outcome. *Reg Anesth Pain Med*. 2005;30:150–162.

Farny J, Drolet P, Girard M. Anatomy of the posterior approach to the lumbar plexus block. *Can J Anaesth*. 1994;41:480–485.

Kirchmair L, Entner T, Kapral S, Mitterschiffthaler G. Ultrasound guidance for the psoas compartment block: an imaging study. *Anesth Analg*. 2002;94:706–710.

Chapter 15

Gray AT, Collins AB, Schafhalter-Zoppoth I. An introduction to femoral nerve and associated lumbar plexus nerve blocks under ultrasonic guidance. *Techniques Reg Anesth Pain Manage*. 2004;8:155–163.

Marhofer P, Nasel C, Sitzwohl C, Kapral S. Magnetic resonance imaging of the distribution of local anesthetic during the three-in-one block. *Anesth Analg*. 2000;90:119–124.

Nielsen KC, Klein SM, Steele SM. Femoral nerve blocks. *Techniques Reg Anesth Pain Manage*. 2003;7:8–17.

Vloka JD, Hadzic A, Drobnik L, Ernest A, Reiss W, Thys DM. Anatomical landmarks for femoral nerve block: a comparison of four needle insertion sites. *Anesth Analg*. 1999;89:1467–1470.

Chapter 16

Bouaziz H, Vial F, Jochum D, et al. An evaluation of the cutaneous distribution after obturator nerve block. *Anesth Analg*. 2002;94:445–449.

Capdevila X, Biboulet P, Bouregba M, Barthelet Y, Rubenovitch J, d'Athis F. Comparison of the three-in-one and fascia iliaca compartment blocks in adults: clinical and radiographic analysis. *Anesth Analg*. 1998;86:1039–1044.

Hopkins PM, Ellis FR, Halsall PJ. Evaluation of local anaesthetic blockade of the lateral femoral cutaneous nerve. *Anaesthesia*. 1991;46:95–96.

Chapter 17

Bollini CA, Moreno M. Sciatic nerve block. *Techniques Reg Anesth Pain Manage*. 2006;10:163–172.

Chan VW, Nova H, Abbas S, McCartney CJ, Perlas A, Xu DQ. Ultrasound examination and localization of the sciatic nerve: a volunteer study. *Anesthesiology*. 2006;104:309–314.

Karmakar MK, Kwok WH, Ho AM, Tsang K, Chui PT, Gin T. Ultrasound-guided sciatic nerve block: description of a new approach at the subgluteal space. *Br J Anaesth*. 2007;98:390–395.

Chapter 18

Chelly JE, Delaunay L. A new anterior approach to the sciatic nerve block. *Anesthesiology*. 1999;91:1655–1660.

Ericksen ML, Swenson JD, Pace NL. The anatomic relationship of the sciatic nerve to the lesser trochanter: implications for anterior sciatic nerve block. *Anesth Analg*. 2002;95:1071–1074.

Vloka JD, Hadzic A, April E, Thys DM. Anterior approach to the sciatic nerve block: the effects of leg rotation. *Anesth Analg*. 2001;92:460–462.

Chapter 19

Domingo-Triado V, Selfa S, Martinez F, et al. Ultrasound guidance for lateral midfemoral sciatic nerve block: a prospective, comparative, randomized study. *Anesth Analg*. 2007;104:1270–1274.

Hadzic A, Vloka JD. A comparison of the posterior versus lateral approaches to the block of the sciatic nerve in the popliteal fossa. *Anesthesiology*. 1998;88:1480–1486.

Taboada MM, Alvarez J, Cortes J, Rodriguez J, Atanassoff PG. Lateral approach to the sciatic nerve block in the popliteal fossa: correlation between evoked motor response and sensory block. *Reg Anesth Pain Med*. 2003;28:450–455.

Chapter 20

Benzon HT, Kim C, Benzon HP, et al. Correlation between evoked motor response of the sciatic nerve and sensory blockade. *Anesthesiology*. 1997;87:547–552.

Sinha A, Chan VW. Ultrasound imaging for popliteal sciatic nerve block. *Reg Anesth Pain Med*. 2004;29:130–134.

Vloka JD, Hadzic A, April E, Thys DM. The division of the sciatic nerve in the popliteal fossa: anatomical implications for popliteal nerve blockade. *Anesth Analg*. 2001;92:215–217.

Chapter 21

Benzon HT, Sharma S, Calimaran A. Comparison of the different approaches to saphenous nerve block. *Anesthesiology*. 2005;102:633–638.

De Mey JC, Deruyck LJ, Cammu G, De Baerdemaeker LE, Mortier EP. A paravenous approach for the saphenous nerve block. *Reg Anesth Pain Med*. 2001;26:504–506.

van der WM, Lang SA, Yip RW. Transsartorial approach for saphenous nerve block. *Can J Anaesth*. 1993;40:542–546.

Chapter 22

Myerson MS, Ruland CM, Allon SM. Regional anesthesia for foot and ankle surgery. *Foot Ankle*. 1992;13:282–288.

Schurman DJ. Ankle-block anesthesia for foot surgery. *Anesthesiology*. 1976;44:348–352.

Chapter 23

Colbern E. The Bier block for intravenous regional anesthesia: technic and literature review. *Anesth Analg*. 1970;49:935–940.

Hilgenhurst G. The Bier block after 80 years: a historical review. *Reg Anesth*. 1990;15:2–5.

Chapter 24

Liu SS, Salinas FV. Continuous plexus and peripheral nerve blocks for postoperative analgesia. *Anesth Analg*. 2003;96:263–272.

Chapter 25

Brull R, McCartney CJ, Chan VW, El-Beheiry H. Neurological complications after regional anesthesia: contemporary estimates of risk. *Anesth Analg*. 2007;104:965–974.

Buckenmaier CC III, Shields CH, Auton AA, et al. Continuous peripheral nerve block in combat casualties receiving low-molecular weight heparin. *Br J Anaesth*. 2006;97:874–877.

Ho M, Ganapathy S. Is peripheral/plexus block safe in the anticoagulated patient? *Techniques Reg Anesth Pain Manag*. 2006;10:56–61.

Horlocker TT, Wedel DJ, Benzon H, et al. Regional anesthesia in the anticoagulated patient: defining the risks (the second ASRA Consensus Conference on Neuraxial Anesthesia and Anticoagulation). *Reg Anesth Pain Med*. 2003;28:172–197.

Chapter 26

Buckenmaier CC, McKnight GM, Winkley JV, et al. Continuous peripheral nerve block for battlefield anesthesia and evacuation. *Reg Anesth Pain Med*. 2005;30:202–205.

Shorten G, Carr DB, Harmon D, Puig MM, Browne J. *Postoperative Pain Management*. Philadelphia, Pa: Saunders Elsevier; 2006.

Chapter 27

Apfelbaum JL, Chen C, Mehta SS, Gan TJ. Postoperative pain experience: results from a national survey suggest postoperative pain continues to be undermanaged. *Anesth Analg*. 2003;97:534–540.

Perkins FM, Kehlet H. Chronic pain as an outcome of surgery. A review of predictive factors. *Anesthesiology*. 2000;93:1123–1133.

Rathmell JP, Wu CL, Sinatra RS, et al. Acute post-surgical pain management: a critical appraisal of current practice, December 2–4, 2005. *Reg Anesth Pain Med*. 2006;31:1–42.

Chapter 28

Cohen SP, Griffith S, Larkin TM, Villena F, Larkin R. Presentation, diagnoses, mechanisms of injury, and treatment of soldiers injured in Operation Iraqi Freedom: an epidemiological study conducted at two military pain management centers. *Anesth Analg*. 2005;101:1098–1103.

White RL, Cohen SP. Return-to-duty rates among coalition forces treated in a forward-deployed pain treatment center: a prospective observational study. *Anesthesiology*. 2007;107:1003–1008.

Chapter 29

Warren J, Fromm RE, Orr RA, Rotello RC, Horst HM. American College of Critical Care Medicine: guidelines for the inter- and intrahospital transport of critically ill patients. *Crit Care Med*. 2004;32:256–262.

Chapter 30

Dalens B, Vanneuville G, Tanguy A. A new parascapular approach to the brachial plexus in children: comparison with the supraclavicular approach.

Anesth Analg. 1987;66:1264–1271.

Puncuh F, Lampugnani E, Kokki H. Use of spinal anaesthesia in paediatric patients: a single centre experience with 1132 cases. *Paediatr Anaesth.* 2004;14:564–567.

Schuepfer G, Johr M. Psoas compartment block in children: Part I—description of the technique. *Paediatr Anaesth.* 2005;15:461–464.

Chapter 31

Frenette L. The acute pain service. *Crit Care Clin.* 1999;15:143–150.

Ready LB. Acute pain: lessons learned from 25,000 patients. *Reg Anesth Pain Med.* 1999;24:499–505.

Werner MU, Soholm L, Rotboll-Nielsen P, Kehlet H. Does an acute pain service improve postoperative outcome? *Anesth Analg.* 2002;95:1361–1372.

Chapter 32

Han J. Physiology of acupuncture: review of thirty years of research. *J Alter Compl Med.* 1997;3:s-101–s-108.

Kaptchuk TJ. Acupuncture: theory, efficacy, and practice. *Ann Intern Med.* 2002;136:374–383.

Niemtzow RC. Battlefield acupuncture. *Medical Acupuncture.* 2007;19(4):225–228. http://www.liebertpub.com/publication.aspx?pub_id=233. Accessed July 1, 2008.

ABBREVIATIONS AND ACRONYMS

AAG: α_1 -acid glycoprotein	MRI: magnetic resonance imaging
ACLS: advanced cardiac life support	MRN: magnetic resonance neurography
AE: aeromedical evacuation	ms: millisecond
AES: aeromedical evacuation system	NBI: nonbattle injury
AMBUS: ambulance bus	NMDA: N-methyl D-aspartate
APS: acute pain service	NSAID: nonsteroidal antiinflammatory drug
ASIS: anterior superior iliac spine	PCA: patient-controlled analgesia
ASP: semipermanent needle (French)	pK: dissociation constant
ATPase: adenosine triphosphatase	PNB: peripheral nerve block
CASEVAC: casualty evacuation	PNS: peripheral nerve stimulator
CCATT: critical care air transport team	PSIS: posterior superior iliac spine
COX: cyclooxygenase	PVB: paravertebral nerve block
CPNB: continuous peripheral nerve block	RATS: Regional Anesthesia Tracking System
CSH: combat support hospital	PSIS: posterior superior iliac spine
CT: computed tomography	SCM: sternocleidomastoid muscle
EMG: electromyography	SH: sacral hiatus
ESI: epidural steroid injection	SI: sacroiliac
FVC: forced vital capacity	STRATEVAC: strategic evacuation
IV: intravenous	TACEVAC: tactical evacuation
LBP: low back pain	TB: trochanteric bursitis
LFC: lateral femoral cutaneous	TFESI: transforaminal epidural steroid injection
MARAA: Military Advanced Regional Anesthesia and Analgesia	USAF: US Air Force
MEDEVAC: medical evacuation	WRAMC: Walter Reed Army Medical Center